

Research Paper



Insights into mental health, lifestyle patterns, and academic attitudes among medical students: a cross-sectional analysis

Dr. Muhammad Imamuzzaman^{1*}, Dr. Kanij Fatema Mukta², Md. Kasif Akhter³,
Dr. Asm Shahariar Ahmed⁴

¹M.B.B.S\MPH (Community Medicine) \MPH (Epidemiology) \Resident, Department of Orthopaedics and Traumatology, Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan.

²M.B.B.S \ Resident, Department of Dermato-venerology, Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan.

³M.B.B.S(last semester)\ Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan.

⁴M.B.B.S \FIPM(India)\ RMC Hospital & Diagnostic Complex LTD, Dhaka, Bangladesh.

Article Info

Article History:

Received: 03 May 2024

Revised: 11 June 2024

Accepted: 17 June 2024

Published: 02 August 2024

Keywords:

Medical Students


Psychological Distress

Mental Health Issues

General Anxiety Disorder

Emotional Exhaustion

Lifestyle Patterns



ABSTRACT

Background: Medical students are more likely to experience mental health issues due to high academic pressure. Therefore, it is essential to have updated information regarding their mental health status.

Methods: A descriptive cross-sectional study was conducted among 406 randomly selected medical students. Informed verbal consent was obtained from all participants. Data were collected through face-to-face interviews, ensuring confidentiality and the right to withdraw at any time. Quality control measures were maintained throughout the study. Data analysis was performed using SPSS version 25.0, and results were presented using descriptive and inferential statistics.

Results: The participants were aged between 18 and 27 years, with a mean age of 22.7±1.8 years. The majority were Muslim (61.3%) and single (85.9%). Psychological distress was observed in 52.7% of students. A significant proportion (72.9%) had a positive attitude toward medical education, although 54.9% felt their current studies were insufficient. Most students (90.7%) reported normal mental health before admission. Nearly half (47.1%) had no generalized anxiety disorder, while only 6.7% required mental health services. Additionally, 63.1% reported good food quality consumption, 45.1% were non-smokers, and 23.6% had a sedentary lifestyle. A majority (69.9%) maintained good relationships with their parents, and 36.5% reported low levels of emotional exhaustion.

Conclusion: Medical students experience considerable psychological distress despite having good mental health prior to admission. While many maintain healthy lifestyles and positive relationships, a notable proportion still require mental health support. These findings highlight the need for targeted mental health interventions among medical students.

Corresponding Author:

Dr. Muhammad Imamuzzaman

M.B.B.S\MPH (Community Medicine) \MPH (Epidemiology)\ Resident, Department of Orthopaedics and Traumatology, Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan.

Email: muhammadimamuzzamanreaz@gmail.com

Copyright © 2024 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

The life of medical students is quite stressful because of their academic journey. Studies show that medical students are especially prone to experiencing challenges with their mental well-being, facing higher rates of mental illness and psychological distress compared to the general population [1]. Medical students and physicians usually face higher rates of mental health problems which include issues like anxiety, depression and breakdown compared to non-medical students. These challenges not only affect personal well-being but also impact care for the patients through decreased empathy and unprofessional behaviours [2]. Various studies have investigated factors that contribute to the mental health challenges faced by medical students. These study reports mentioned extreme workload, professional competition, reduced sleep duration, lack of support from society, and patient suffering and death. These factors can significantly impact students' stress levels and overall mental well-being [3]. Medical education's specific pathogenetic factors lead to higher stress-related illnesses like depression and burnout in students compared to their peers in other fields or employment [4]. Medical students often avoid seeking mental health care despite the clear need for it. Concerns about privacy, stigma, and potential negative impacts on their careers deter many from seeking help. This avoidance due to stigma-related barriers may lead to increased distress and drained among future practising physicians [5]. Medical school and early career stages for doctors are highly stressful, leading to risks like anxiety, burnout, depression, and alcohol abuse. Even before graduation, students experience significant distress, surpassing peers in their age group. These challenges stem from the combined pressures of academics and personal life [6]. In mental health research, this shift would move away from simply assessing the presence or absence of psychological distress to a more comprehensive evaluation. This broader approach considers not only distress but also factors like psychological well-being and coping strategies to determine a person's mental [7]. This is very important to evaluate the mental health condition of each individual especially who are in the field of medical science because of their tremendous workload. Because of such requirements, current research was conducted.

2. RELATED WORK

Medical students are especially susceptible to experiencing stress, anxiety, and depression. Recent research has highlighted significant rates of mental health issues among medical trainees globally. Recent research involving 129,123 medical students from 47 countries found high rates of stress-related issues. Around 27.2% showed symptoms of depression, with 11.1% reporting suicidal thoughts [8]. Research has consistently highlighted poor mental health among medical students in various Asian countries such as India, Pakistan, Iran, Malaysia, China, and Saudi Arabia. This includes issues like serious contemplation of dropping out of medical school, substance abuse, burnout, and suicidal thoughts, emphasizing the significant challenges faced by students in these regions [9]. Medical education is often seen as stressful, leading to issues like anxiety, depression, and burnout due to heavy workloads, frequent testing, and

increasing patient responsibilities. Longitudinal studies indicate a worsening trend in mental health and well-being during medical training [10].

3. METHODOLOGY

A descriptive cross-sectional study was conducted among 406 randomly selected medical students in Bishkek, Kyrgyzstan. Participants were chosen through random selection methods within the student population. Before data collection, informed verbal consent was obtained from each participant, ensuring ethical standards were met. Data collection was carried out by the principal investigator using a pretested interview schedule through face-to-face interviews, ensuring privacy and data quality. Quality control measures included checking questionnaires for completeness, consistency, correctness, and discrepancies. Confidentiality of collected data was maintained throughout the study. Participants were informed of their right to withdraw from the study at any time without providing a reason. No physical, social, or psychological risks were associated with the study procedure. Data analysis was performed using SPSS version 25.0, employing descriptive statistics such as frequency, percentage, mean, and standard deviation. Inferential statistics, specifically the Chi-Square test, were used where appropriate. Results were presented using tables and diagrams for clarity and understanding.

4. RESULTS AND DISCUSSION

The age range of the respondents was 18 to 27 years, while a majority of 288(70.9%) of the students were in the age group 20 to 24 years with a mean age of 22.7 ± 1.8 years. In terms of gender distribution, 212(52.2%) were male, while 194(47.8%) were female. Of 406 medical students, the religion of 249(61.3%) were muslim while 151(37.2%) were hindu and 6(1.5%) were Buddhist. Marital status indicated that a significant 349(85.9%) were single, while 57(14.1%) were married. Regarding type of family, it was revealed that 284(69.9%) had nuclear family and 122(30.1%) had Extended Family. Of the students, 354 (87.2%) of students opined that their family's economic status was good while 52(12.8%) had poor economic status. The majority 233(57.4%) of students were from private medical colleges and 173(42.6%) from state medical.

A significant number 214(52.7%) had psychological distress while 192(47.3%) had no psychological distress. Near three-fourth 296(72.9%) had a positive attitude towards choosing medical education and 110(27.1%) had no opinion or negative attitude about it. The majority 223(54.9%) respondents revealed that their study was not sufficient, 97(23.9%) opined sufficient and 86(21.2%) had no opinion about it. Of them, 38(9.3%) had mental health issues before admission while 368(90.7%) had normal mental health. Only 27(6.7%) had received mental health services and nearly cent 379(93.3%) were not required to have mental health services. Around half 191(47.1%) of respondents had no general anxiety disorder, 166(40.9%) had mild while 28(6.9%) had moderate and 21(5.1%) had severe anxiety disorder [Table 1](#).

Table 1. Respondents by General Anxiety Disorder (GAD)

General Anxiety Disorder (GAD)	Frequency	Percentage
No GAD	191	47.1
Mild GAD	166	40.9
Moderate GAD	28	6.9
Severe GAD	21	5.1
Total	406	100.00

Regarding current mental health disturbance, 191(47.1%) had no mental health condition, 76(18.7%) had psychological distress, 36(8.9%) had social dysfunction and 103(25.3%) had both psychological distress and social dysfunction [Figure 1](#).




Figure 1. Respondents by Current Mental Health Disturbance

Near two-thirds, 256(63.1%) had good food quality and 150(36.9%) had average quality food. Among the student, 168(41.4%) consume unhealthy diet, 131(32.3%) had average food and 107(26.3%) consume healthy food. Majority of respondents 183(45.1%) were non-smokers, 96(23.6%) were occasional smoker and 127(31.3%) were daily smokers Table 2.

Table 2. Respondents by Smoking Status

Smoking Status	Frequency	Percentage
Never	183	45.1
Occasional	96	23.6
Daily	127	31.3
Total	406	100.00

Around one-fourths of medical students, 96(23.6%) had sedentary lifestyle, 125(30.8%) had moderate physical activity and 185(45.6%) had active lifestyle. A significant number 284(69.9%) had good relationships with their parents, 94(23.2%) had average relations and 28(6.9%) had poor relations with their parents. More than one-third 148(36.5%) had a low level of emotional exhaustion, 184(45.3%) had moderate and 74(18.2%) had a high level of emotional exhaustion. Regarding the association, it was found that among the medical students, there is a statistically significant association ($p < 0.005$) between psychological distress and physical activity Table 3.

Table 3. Association between Psychological Distress and Physical Activity

Psychological Distress	Physical Activity			Total N (%)	P Value
	Sedentary	Moderate	Active		
Yes	74	109	31	214(100)	<0.005
No	22	16	154	192(100)	
Total	96	125	185	406(100)	

4.1. Discussion

Respondents of current study were between 18 to 24 years with a mean age of 21.3 ± 0.9 years. This is consistent, where 56.1% of the medical students were between 21 to 30 years old [11]. Regarding gender, 212(52.2%) were male, while 194(47.8%) were female. This is inconsistent where 30.55% were male and 69.45 were female which may be due to the small sample size, and purposive selection of the study group

[12]. More than two-fourths of the current study were 349(85.9%) single which is corroborated where the majority 99.1% were single [13]. More than two-thirds 284(69.9%) of respondents had nuclear family. This is vindicated by where 86.1% had a nuclear family [14]. In this study, 354(87.2%) of students but 52(12.8%) families had poor economic conditions. This is vindicated by findings of the study conducted by Carletto S and associates where the majority 94.1% had good economic status [15]. In this study, around two-fifths of 173(42.6%) were students of state medical and 233(57.4%) from private medical colleges. This is substantiated by findings of a study conducted by Zila-Velasque JP and team where 35.1% were studying in the state and 64.9% were from private medical colleges [16]. Of the respondents, around half 214(52.7%) were suffering from psychological distress but 192(47.3%) were not. Where they found that the prevalence of psychological distress was 51% [17]. In the present study, 296(72.9%) had a positive attitude towards choosing medical education and 110(27.1%) had no opinion or negative attitude towards it. This is more or less similar to the findings of Carletto S and team where more than two-thirds 67.6% had positive and the rest 32.4% had negative/no opinions about it [15]. Of students, more than half 223(54.9%) opined that their study was sufficient and 97(23.9%) were not satisfied while 86(21.2%) had no opinion about it. Where 26.73% thought their study was sufficient while 52.48% were not but 20.79% were unsure about it. This difference might be attributed to variations in study area and in sample size [18]. The majority 368(90.7%) had normal mental health before medical college admission and only 38(9.3%) had mental health issues during pre-admission time. This is corroborated by, where 96.8% had normal mental health conditions and 3.2% had mental health issues during pre-admission [19]. Among the study respondents, 379(93.3%) of the respondents need not have mental health services and only 27(6.7%) received mental health services. This is corroborated by who found that the majority 86.73% not receive mental health services while only 13.27% need to have such services [20].

Regarding General Anxiety Disorder(GAD) , of respondents 191(47.1%) had no GAD, 166(40.9%) had mild, 28(6.9%) had moderate and only 21(5.1%) had severe disorder. This is substantiated by findings of a study conducted by Christophers B and team, where 42% had no GAD, 38% had mild, 13 had moderate and only 7% had severe GAD [21]. Regarding current mental health disturbance, 191(47.1%) had no mental health condition, 76(18.7%) had psychological distress, 36(8.9%) had social dysfunction and 103(25.3%) had both psychological distress and social dysfunction. This is supported by where 50.9% were currently mentally healthy, 12.8% had psychological distress, 15.9% had social dysfunction but 20.4% opined to both have psychological distress and social dysfunction [22]. A significant number 256(63.1%) had good food quality and 150(36.9%) had the average quality of food. This is vindicated by findings of the study conducted by Kobbaz TM and team, where more than half 55.4% had good quality eating while 44.5% had poor quality [23]. Of students, 168(41.4%) consume unhealthy diet, 131(32.3%) had average food and 107(26.3%) consume healthy food. A study conducted by found nearly similar results where 33.2% consumed unhealthy diets, 51.6% had average diets and only 15.3% consumed a healthy diet [24]. Of respondents, 183(45.1%) were non-smokers, 96(23.6%) were occasional smoker and 127(31.3%) were daily smokers. This is supported by findings of the study conducted where 56.1% never smoked, 19.3% were occasional smokers and nearly one-fourth 24.6% smoke daily [25]. Around half 185(45.6%) had active lifestyle, 125(30.8%) had moderate activity and 96(23.6%) had sedentary lifestyle. This is in line with the study findings of where only 11.8% were inactive, 13.2% were moderately active and 75.0% were active [24]. Regarding relationships with the parents, 284(69.9%) had good relationships with their parents, 94(23.2%) had average relations and 28(6.9%) had poor relations with their parents. This is substantiated where 80.5% had good relations with their parents and 19.5% did not have that much good relationship with their parents [26]. Of the study respondents, 148(36.5%) had a low level of emotional exhaustion, 184(45.3%) had moderate and 74(18.2%) had a high level of emotional exhaustion. This is vindicated by findings of the study conducted by Pitanupong J and team where 14.9% had a low level of emotional exhaustion, 22.0% had moderate, 62.8% had a high level of emotional exhaustion and 0.3% had no answer about it [27].

5. CONCLUSION

Medical students were mostly Muslim single males from nuclear families with good economic status studying in private medical colleges. Respondents were suffering from psychological distress. Students had positive attitudes about medical education and their pre-admission mental health was good but not satisfied with their current study. Students were suffering from mental health conditions but little had received mental health services. Students consume healthy and good quality food while most of them are non-smokers and have an active lifestyle. Respondents had a good relationship with their parents and suffered from low levels of emotional exhaustion.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author Contributions Statement

Name of Author	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Dr. Muhammad Imamuzzaman	✓	✓	✓	✓	✓	✓			✓	✓	✓	✓	✓	
Dr. Kanij Fatema Mukta		✓		✓		✓		✓	✓			✓		✓
Md. Kasif Akhter	✓	✓		✓	✓		✓			✓		✓	✓	
Dr. Asm Shahariar Ahmed	✓		✓		✓		✓	✓		✓	✓		✓	✓

C: Conceptualization

M: Methodology

So: Software

Va: Validation

Fo: Formal analysis

I: Investigation

R: Resources

D: Data Curation

O: Writing- Original Draft

E: Writing- Review & Editing

Vi: Visualization

Su: Supervision

P: Project administration

Fu: Funding acquisition

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study, and their voluntary consent was obtained prior to data collection.

Ethical Approval

The study was conducted in compliance with the ethical principles outlined in the Declaration of Helsinki and approved by the relevant institutional authorities.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES


- [1] Z. Lyons, H. Wilcox, L. Leung, and O. Dearsley, 'COVID-19 and the mental well-being of Australian medical students: impact, concerns and coping strategies used', *Australas. Psychiatry*, vol. 28, no. 6, pp. 649-652, Dec. 2020. doi.org/10.1177/1039856220947945




- [2] B. Van Remortel, E. Dolan, D. Cipriano, and P. McBride, 'Medical Student Wellness in Wisconsin: Current Trends and Future Directions', *WMJ*, vol. 117, no. 5, pp. 211-213, Dec. 2018.
- [3] T. Kötter, Y. Tautphäus, M. Scherer, and E. Voltmer, 'Health-promoting factors in medical students and students of science, technology, engineering, and mathematics: design and baseline results of a comparative longitudinal study', *BMC Med. Educ.*, vol. 14, no. 1, p. 134, July 2014. doi.org/10.1186/1472-6920-14-134
- [4] L. N. Dyrbye, M. R. Thomas, and T. D. Shanafelt, 'Systematic review of depression, anxiety, and other indicators of psychological distress among U.S. and Canadian medical students', *Acad Med*, vol. 81, pp. 354-373, 2006. doi.org/10.1097/00001888-200604000-00009
- [5] J. A. Lavergne and M. L. Kennedy, 'Telepsychiatry and medical students: A promising mental health treatment for medical student use both personally and professionally', *Curr. Psychiatry Rep.*, vol. 23, no. 6, p. 31, Apr. 2021. doi.org/10.1007/s11920-021-01248-6
- [6] V. Yogeswaran and C. El Morr, 'Effectiveness of online mindfulness interventions on medical students' mental health: a systematic review', *BMC Public Health*, vol. 21, no. 1, p. 2293, Dec. 2021. doi.org/10.1186/s12889-021-12341-z
- [7] I. van Dijk, P. L. B. J. Lucassen, C. van Weel, and A. E. M. Speckens, 'A cross-sectional examination of psychological distress, positive mental health and their predictors in medical students in their clinical clerkships', *BMC Med. Educ.*, vol. 17, no. 1, Dec. 2017. doi.org/10.1186/s12909-017-1035-8
- [8] A. Damião Neto, A. L. G. Lucchetti, O. da Silva Ezequiel, and G. Lucchetti, 'Effects of a required large-group mindfulness meditation course on first-year medical students' mental health and quality of life: A randomized controlled trial', *J. Gen. Intern. Med.*, vol. 35, no. 3, pp. 672-678, Mar. 2020. doi.org/10.1007/s11606-019-05284-0
- [9] A. Adhikari, A. Dutta, S. Sapkota, A. Chapagain, A. Aryal, and A. Pradhan, 'Prevalence of poor mental health among medical students in Nepal: a cross-sectional study', *BMC Med. Educ.*, vol. 17, no. 1, Dec. 2017. doi.org/10.1186/s12909-017-1083-0
- [10] E. Voltmer, K. Obst, and T. Kötter, 'Study-related behavior patterns of medical students compared to students of science, technology, engineering and mathematics (STEM): a three-year longitudinal study', *BMC Med. Educ.*, vol. 19, no. 1, p. 262, July 2019. doi.org/10.1186/s12909-019-1696-6
- [11] J. Baj-Korpak, K. Zaworski, E. Szymczuk, and A. Shpakou, 'Physical activity and mental health of medical students from Poland and Belarus-countries with different restrictive approaches during the COVID-19 pandemic', *Int. J. Environ. Res. Public Health*, vol. 19, no. 21, p. 13994, Oct. 2022. doi.org/10.3390/ijerph192113994
- [12] P. Xiong et al., 'Factors influencing mental health among Chinese medical and non-medical students in the early stage of the COVID-19 pandemic', *Front. Public Health*, vol. 9, p. 603331, May 2021. doi.org/10.3389/fpubh.2021.603331
- [13] A. Hanani, M. Badrasawi, S. Zidan, and M. Hunjul, 'Effect of cognitive behavioral therapy program on mental health status among medical student in Palestine during COVID pandemic', *BMC Psychiatry*, vol. 22, no. 1, p. 310, May 2022. doi.org/10.1186/s12888-022-03915-1
- [14] B. Hu, Y. Wang, H. T. Zhou, M. Li, and L. Zheng, 'A study on the impact of mental health problems on the academic buoyancy of medical students and the mechanisms', *Front. Public Health*, vol. 11, p. 1229343, 2023. doi.org/10.3389/fpubh.2023.1229343
- [15] S. Carletto et al., 'The impact of COVID-19 on mental health in medical students: A cross-sectional survey study in Italy', *Psychol. Rep.*, vol. 127, no. 2, pp. 620-648, Apr. 2024. doi.org/10.1177/00332941221127632
- [16] J. P. Zila-Velasque et al., 'Sociodemographic and educational factors associated with mental health disorders in medical students of clinical years: A multicenter study in Peru', *PLoS One*, vol. 18, no. 6, p. e0286338, June 2023. doi.org/10.1371/journal.pone.0286338
- [17] P. K. Sahu, B. S. Nayak, V. Rodrigues, and S. Umakanthan, 'Prevalence of psychological distress among undergraduate medical students: A cross-sectional study', *Int. J. Appl. Basic Med. Res.*, vol. 10, no. 4, pp. 270-275, Oct. 2020. doi.org/10.4103/ijabmr.IJABMR_100_19

- [18] M. Karabacak et al., 'Final-year medical student mental wellness during preparation for the examination for specialty in Turkey: a cross-sectional survey study', *BMC Med. Educ.*, vol. 23, no. 1, p. 79, Feb. 2023. doi.org/10.1186/s12909-023-04063-0
- [19] S. Wainipitapong and M. Chiddaycha, 'Assessment of dropout rates in the preclinical years and contributing factors: a study on one Thai medical school', *BMC Med. Educ.*, vol. 22, no. 1, p. 461, June 2022. doi.org/10.1186/s12909-022-03527-z
- [20] R. B. Kihumuro et al., 'Knowledge, attitude and perceptions of medical students towards mental health in a university in Uganda', *BMC Med. Educ.*, vol. 22, no. 1, p. 730, Oct. 2022. doi.org/10.1186/s12909-022-03774-0
- [21] B. Christophers, E. Nieblas-Bedolla, J. S. Gordon-Elliott, Y. Kang, K. Holcomb, and M. K. Frey, 'Mental health of US medical students during the COVID-19 pandemic', *J. Gen. Intern. Med.*, vol. 36, no. 10, pp. 3295-3297, Oct. 2021. doi.org/10.1007/s11606-021-07059-y
- [22] T. Wiguna, V. J. Dirjayanto, Z. S. Maharani, E. G. Faisal, S. D. Teh, and E. Kinzie, 'Mental health disturbance in preclinical medical students and its association with screen time, sleep quality, and depression during the COVID-19 pandemic', *BMC Psychiatry*, vol. 24, no. 1, p. 85, Jan. 2022. doi.org/10.1186/s12888-024-05512-w
- [23] T. M. Kobbaz et al., 'The lifestyle of Brazilian medical students: What changed and how it protected their emotional wellbeing during the COVID-19 pandemic', *Aust. J. Gen. Pract.*, vol. 50, no. 9, pp. 668-672, Sept. 2021. doi.org/10.31128/AJGP-03-21-5886
- [24] A. Terebessy, E. Czeglédi, B. C. Balla, F. Horváth, and P. Balázs, 'Medical students' health behaviour and self-reported mental health status by their country of origin: a cross-sectional study', *BMC Psychiatry*, vol. 16, no. 1, Dec. 2016. doi.org/10.1186/s12888-016-0884-8
- [25] C. A. Popescu et al., 'Evolving mental health dynamics among medical students amid COVID-19: A comparative analysis of stress, depression, and alcohol use among medical students', *Medicina (Kaunas)*, vol. 59, no. 10, p. 1854, Oct. 2023. doi.org/10.3390/medicina59101854
- [26] Z. Gui, L. Sun, and C. Zhou, 'Self-reported sleep quality and mental health mediate the relationship between chronic diseases and suicidal ideation among Chinese medical students', *Sci. Rep.*, vol. 12, no. 1, p. 18835, Nov. 2022. doi.org/10.1038/s41598-022-23207-8
- [27] Pitanupong J, Sathaporn K, Ittasakul P, Karawekpanyawong N. Relationship of mental health and burnout with empathy among medical students in Thailand: A multicenter cross-sectional study. *PLoS One*. 2023 Jan 5;18(1):e0279564. doi: 10.1371/journal.pone.e0279564. PMID: 36602955; PMCID: PMC9815634. doi.org/10.1371/journal.pone.0279564

How to Cite: Dr. Muhammad Imamuzzaman, Dr. Kanij Fatema Mukta, Md. Kasif Akhter, Dr. Asm Shahariar Ahmed. (2024). Insights into mental health, lifestyle patterns, and academic attitudes among medical students: a cross-sectional analysis. *Journal of Prevention, Diagnosis and Management of Human Diseases (JPDMHD)*, 4(2), 19-27. <https://doi.org/10.55529/jpdmhd.44.19.27>

BIOGRAPHIES OF AUTHORS

	<p>Dr. Muhammad Imamuzzaman, is an MBBS graduate with dual MPH degrees in Community Medicine and Epidemiology. He is currently a Resident in the Department of Orthopaedics and Traumatology at Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan. His academic interests include public health research, epidemiology, mental health, and evidence-based clinical practice. He has actively participated in cross-sectional and community-based studies focusing on student health, lifestyle patterns, and preventive medicine. His work integrates clinical training with population health research to improve healthcare outcomes. Email: muhammadimamuzzamanreaz@gmail.com</p>
---	--

	<p>Dr. Kanij Fatema Mukta, is an MBBS graduate and currently serves as a Resident in the Department of Dermato-venerology at Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan. Her professional interests include dermatological disorders, sexually transmitted infections, and community-based health research. She has been involved in academic studies addressing mental health and lifestyle behaviors among medical students. She is committed to integrating clinical dermatology with preventive and public health approaches to enhance patient care and awareness.</p>
	<p>Md. Kasif Akhter, is a final-semester MBBS student at Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan. He has a keen interest in clinical research, medical education, and student mental health. Throughout his academic journey, he has actively contributed to research projects focusing on psychological well-being and lifestyle factors among medical students. He aims to pursue a career that combines clinical excellence with research and public health initiatives to promote holistic healthcare delivery.</p>
	<p>Dr. Asm Shahariar Ahmed, is an MBBS graduate with FIPM (India) and is currently affiliated with RMC Hospital & Diagnostic Complex Ltd., Dhaka, Bangladesh. His professional interests include clinical medicine, pain management, and hospital-based healthcare services. He has contributed to collaborative research projects focusing on mental health and lifestyle patterns among medical students. He is dedicated to advancing patient-centered care through clinical expertise and continuous academic involvement.</p>